

Immobilization of recombinant *Escherichia coli* on multi-walled carbon nanotubes for xylitol production

Noor Hidayah Abd Rahman^a, Jamaliah Md. Jahim^b, Mimi Sakinah Abdul Munaim^c, Roshanida A. Rahman^a, Siti Fatimah Zaharah Fuzi^d, Rosli Md. Illias^{a,e}

^aSchool of Chemical and Energy Engineering, Faculty of Engineering, Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

^bFaculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

^cFaculty of Engineering Technology, Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300 Gambang, Kuantan, Pahang, Malaysia

^dFaculty of Science, Technology and Human Development, Universiti Tun Hussein Onn Malaysia, 86400 Batu Pahat, Johor, Malaysia

^eInstitute of Bioproduct Development, Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

ABSTRACT

E. coli has been engineered to produce xylitol, but the production faces bottlenecks in terms of production yield and cell viability. In this study, recombinant *E. coli* (r*E. coli*) was immobilized on untreated and treated multiwalled carbon nanotubes (MWCNTs) for xylitol production. The immobilized r*E. coli* on untreated MWCNTs gave the highest xylitol production (5.47 g L^{-1}) and a productivity of $0.22 \text{ g L}^{-1} \text{ h}^{-1}$. The doubling time for the immobilized cells increased up to 20.40 h and was higher than that of free cells (3.67 h). Cell lysis of the immobilized cells was reduced by up to 73 %, and plasmid stability improved by up to 17 % compared to those of free cells. Xylitol production using the optimum parameters (pH 7.4, 0.005 mM and 29 °C) achieved a xylitol production and productivity of 6.33 g L^{-1} and $0.26 \text{ g L}^{-1} \text{ h}^{-1}$, respectively. A seven-cycle repeated batch fermentation was carried out for up to 168 h, which showed maximum xylitol production of 7.36 g L^{-1} during the third cycle. Hence, this new adsorption immobilization system using MWCNTs is an alternative to improve the production of xylitol.

KEYWORDS

Xylitol; Cell immobilization; *Escherichia coli*; multi-walled carbon nanotubes; Shake flask fermentation

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the assistance of the Ministry of Higher Education of Malaysia under the Long-Term Research Grant Scheme (LRGS/2013/UKM-UKM/PT/01) and the Universiti Teknologi Malaysia (UTM).